

PROCESS FOR REDUCING KETOCARBOXYLIC ESTERS

5

## BACKGROUND OF THE INVENTION

Field of the Invention: The present invention relates to a process for preparing enantiomerically enriched alpha- and beta-hydroxycarboxylic esters from the corresponding ketocarboxylic esters and also relates to immobilized transition metal complexes usable therefor.

Brief Description of the Prior Art: Enantiomerically enriched alpha- and beta-hydroxycarboxylic esters are valuable reagents for optical resolution and important intermediates in the preparation of pharmaceuticals and agrochemicals.

10 Customarily, enantiomerically enriched alpha- and beta-hydroxycarboxylic esters are obtained by the process of catalytically hydrogenating the corresponding alpha- and beta-ketocarboxylic esters, usually using transition metal complexes having chiral phosphines as ligands as catalysts (see, for example, Genet et al., Tetrahedron, Asymmetry, 1994, 5(4), 675-690). A disadvantage of chiral  
15 phosphines is their high cost and oxidation sensitivity, which is why they are used on the industrial scale predominantly in homogeneous processes, if at all.

Alternatively, processes using platinum or nickel catalysts modified by quinchonaalkaloids or tartaric acid derivatives are known (T. Mallat et al., Fine  
20 Chemicals through Heterogeneous Catalysis, Wiley-VCH, 2001, p. 449 ff).

Also, Ferrand et al. (Tetrahedron: Asymmetry, 13, 2002, pp. 1379 to 1384) describe the use of rhodium, ruthenium and iridium complexes with chiral diamines for the hydrogenation of ketoesters.

25

A common disadvantage of all these processes is that they allow at best a moderate enantiomeric excess.


"Express Mail" mailing label number EU942980572US

Date of Deposit February 3, 2004

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the ~~Assistant~~ Commissioner of Patents and Trademarks, ~~Washington, D.C. 20231~~ ALEXANDRIA, VA

Sara Sue Riley

(Name of person mailing paper or fee)


  
Signature of person mailing paper or fee

There was therefore a need to provide catalysts which make possible high yields and enantioselectivities in particular in a process for preparing enantiomerically enriched alpha- and beta-hydroxycarboxylic esters.


5

# SUMMARY OF THE INVENTION

In accordance with the foregoing, the present invention encompasses compounds of the formula (I)



10 where



is an enantiomerically enriched chiral nitrogen compound,

Linker is a radical which is bonded both covalently to the enantiomerically enriched chiral nitrogen compound and to the support,

15

Support is a micro-, meso- or macroporous support material,

(M<sup>m+</sup>) is a metal having valency m

20 L is an anionic or uncharged ligand

n is one, two, three or four

(An  $q^-$ ) is an anion having valency  $q$  and

$p$  is  $(m - \text{number of anionic ligands } L)/q$ .

- 5 For the purposes of the invention, enantiomerically enriched compounds are enantiomerically pure compounds or mixtures of enantiomers of a compound in which one enantiomer is present in an enantiomeric excess, (also referred to hereinbelow as ee) relative to the other enantiomer. Preferably, this enantiomeric excess is 10 to 100% ee, particularly preferably 90 to 100% ee and very particularly preferably 95 to 100% ee.
- 10

For the purposes of the invention, all radical definitions, parameters and illustrations hereinabove and listed hereinbelow, in general or within areas of preference, i.e. the particular areas and areas of preference, may be combined as desired.

15

#### DETAILED DESCRIPTION OF THE INVENTION

Alkyl, alkoxy, alkylene and alkenylene hereinbelow are each independently a straight-chain, cyclic, branched or unbranched alkyl, alkoxy, alkylene and alkenylene radical respectively, each of which may optionally be further substituted by  $C_1$ - $C_4$ -alkoxy. The same applies to the nonaromatic moiety of an arylalkyl radical.

20

Illustrative but non-limiting examples of these radicals are as follows.  $C_1$ - $C_4$ -Alkyl is, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl and tert-butyl,  $C_1$ - $C_8$ -alkyl is additionally, for example, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, neopentyl, 1-ethylpropyl, cyclohexyl, cyclopentyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-

25

dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl and n-octyl, and C<sub>1</sub>-C<sub>20</sub>-alkyl is further additionally, for example, adamantyl, the isomeric menthyls, n-nonyl, n-decyl and n-dodecyl.

5

C<sub>1</sub>-C<sub>4</sub>-Alkoxy is, for example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy and tert-butoxy, C<sub>1</sub>-C<sub>8</sub>-alkoxy is additionally, for example, n-pentoxo, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, neopentoxo, 1-ethylpropoxy, cyclohexoxy, cyclopentoxo, n-hexoxy and n-octoxy, and C<sub>1</sub>-C<sub>20</sub>-alkoxy is further additionally, for example, adamantoxo, the isomeric menthoxy radicals, n-decoxy and n-dodecoxy.

10

C<sub>1</sub>-C<sub>4</sub>-Alkylene is, for example, methylene, 1,1-ethylene, 1,2-ethylene, 1,1-propylene, 1,3-propylene, 1,4-butylene, and C<sub>1</sub>-C<sub>8</sub>-alkylene is additionally, for example, 1,2-cyclohexylene and 1,2-cyclopentylene.

15

C<sub>2</sub>-C<sub>8</sub>-Alkenylene is, for example, 1,1-ethenylene 2-ethoxy-1,1-ethenylene and 2-methoxy-1,1-ethenylene.

20

**Haloalkyl, haloalkoxy and haloalkylene** are each independently a straight-chain, cyclic, branched or unbranched alkyl radical and alkoxy radical and alkylene radical respectively, each of which is singly, multiply or fully substituted by halogen atoms.

25

For example, C<sub>1</sub>-C<sub>20</sub>-haloalkyl is trifluoromethyl, chloromethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, nonafluorobutyl, heptafluoroisopropyl, perfluorooctyl, perfluorodecyl and perfluorohexadecyl.

**Aryl** is in each case independently a heteroaromatic radical having 5 to 14 framework carbon atoms of which no, one, two or three framework carbon atoms per cycle, but at least one framework carbon atom in the entire molecule, may be substituted by heteroatoms selected from the group of nitrogen, sulphur or oxygen, or and is preferably a carbocyclic aromatic radical having 6 to 14 framework carbon atoms.

Examples of carbocyclic aromatic radicals having 6 to 14 framework carbon atoms are phenyl, biphenyl, naphthyl, phenanthrenyl, anthracenyl or fluorenyl, heteroaromatic radicals having 5 to 14 framework carbon atoms of which no, one, two or three framework carbon atoms per cycle, but at least one framework carbon atom in the entire molecule, may be substituted by heteroatoms selected from the group of nitrogen, sulphur or oxygen are, for example, pyridinyl, oxazolyl, benzofuranyl, dibenzofuranyl or quinolinyl.


The carbocyclic aromatic radical or heteroaromatic radical may also be substituted by up to five identical or different substituents per cycle which are selected, for example, from the group of nitro, cyano, chlorine, fluorine, C<sub>1</sub>-C<sub>12</sub>-alkyl, C<sub>1</sub>-C<sub>12</sub>-haloalkyl, C<sub>1</sub>-C<sub>12</sub>-haloalkoxy, C<sub>1</sub>-C<sub>12</sub>-haloalkylthio, C<sub>1</sub>-C<sub>12</sub>-alkoxy, di(C<sub>1</sub>-C<sub>8</sub>-alkyl)amino or tri(C<sub>1</sub>-C<sub>6</sub>-alkyl)siloxy.


**Arylene** is an aryl radical which has a further bonding site on the aromatic framework and is therefore divalent.

**Arylalkyl** is in each case independently a straight-chain, cyclic, branched or unbranched alkyl radical as defined above which may be singly, multiply or fully substituted by aryl radicals as defined above.

**Arylalkylene** is an arylalkyl radical which has a further bonding site on the aromatic framework and is therefore divalent.

Areas of preference for compounds of the formula (I) are defined hereinbelow:

5  is preferably an enantiomerically enriched chiral nitrogen compound of the formula (II)



where

10

the arrow indicates the bonding point to the linker and

15  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^4$  are each independently hydrogen,  $\text{C}_1$ - $\text{C}_8$ -alkyl,  $\text{C}_5$ - $\text{C}_{15}$ -arylalkyl or  $\text{C}_4$ - $\text{C}_{14}$ -aryl or  $\text{NR}^1\text{R}^2$  as a whole is a cyclic amino radical having a total of 4 to 20 carbon atoms,

$\text{R}^3$  is a divalent radical having 2 to 30 carbon atoms or

20  $\text{R}^3$  and at least one of the radicals  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^4$  together are part of a cyclic amino radical having a total of 4 to 20 carbon atoms.

Preferred compounds of the formula (II) are those in which

25  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^4$  are each independently hydrogen,  $\text{C}_1$ - $\text{C}_8$ -alkyl,  $\text{C}_5$ - $\text{C}_{15}$ -arylalkyl or  $\text{C}_4$ - $\text{C}_{14}$ -aryl or  $\text{NR}^1\text{R}^2$  as a whole is a 5- or 6-membered monocyclic amino

radical which is optionally mono-, di-, tri- or tetrasubstituted on the carbon framework by C<sub>1</sub>-C<sub>4</sub>-alkyl and

5 R<sup>3</sup> is a divalent radical which is selected from the group of C<sub>2</sub>-C<sub>8</sub>-alkylene which may optionally be further mono- or disubstituted by C<sub>4</sub>-C<sub>14</sub>-aryl radicals, C<sub>5</sub>-C<sub>15</sub>-arylalkylene, C<sub>4</sub>-C<sub>14</sub>-arylene or bis(C<sub>4</sub>-C<sub>14</sub>-arylene) or

10 R<sup>3</sup> and one of the radicals R<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> together are part of a 5- or 6-membered monocyclic amino radical which is optionally additionally mono-, di-, tri- or tetrasubstituted on the carbon framework by C<sub>1</sub>-C<sub>4</sub>-alkyl.

Particularly preferred compounds of the formula (II) are those in which

15 R<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> are each independently hydrogen, methyl or ethyl and

R<sup>3</sup> is a divalent radical which is selected from the group of 1,2-bis(C<sub>4</sub>-C<sub>14</sub>-aryl)-1,2-ethylene, 1,2-cyclohexylene, 1,1'-2,2'-bis(C<sub>4</sub>-C<sub>14</sub>-arylene) or

20 R<sup>3</sup> and one of the radicals R<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> together are part of a pyrrolidinyl or piperidinyl radical.

Very particularly preferred compounds of the formula (II) are those which are derived from the following compounds:

25 (1R,2R)-1,2-diphenylethylenediamine, (1S,2S)-1,2-diphenylethylenediamine,  
(1R,2R)-1,2-dimethylethylenediamine, (1S,2S)-1,2-dimethylethylenediamine,  
(1R,2R)-1,2-cyclohexylenediamine, (1S,2S)-1,2-cyclohexylenediamine, (S)-2-aminomethyl-1-ethylpyrrolidine, (R)-2-aminomethyl-1-ethylpyrrolidine, (S)-2-aminomethyl-1-methylpyrrolidine, (R)-2-aminomethyl-1-methylpyrrolidine, (R)-

1,1'-diamino-2,2'-binaphthyl, (S)-1,1'-diamino-2,2'-binaphthyl, (R)-1,1'-diamino-6,6'-dimethoxy-2,2'-biphenyl and (S)-1,1'-diamino-6,6'-dimethoxy-2,2'-biphenyl, and even greater preference is given to (R)-2-aminomethyl-1-ethylpyrrolidine and (S)-2-aminomethyl-1-methylpyrrolidine.

5

**Support** is preferably a micro- or mesoporous support material. The terms micro-, meso- and for that matter macroporous, and the nomenclature of the zeolites are to be interpreted in accordance with IUPAC (McCusker et al. Pure Appl. Chem, vol. 73, No. 2, pp. 381-394, 2001). Examples of suitable support materials include

10 silica gels, or zeolites of the MOR, X, Y, MCM, ZSM the , FAU, MFI, L, BEA, FER, A and SBA type or those of the AlPO, MAIPO and SAPO type, and the zeolites mentioned may optionally be isomorphically substituted. Particular preference is given to mesoporous zeolites, in particular those of the MCM type, for example MCM-41.

15

Preferred **linker-support** combinations are those which are obtainable by initially organically modifying the surface of the support in such a way that it has one or more functionalities after the modification (referred to hereinbelow as **activated support**) and that it is bonded via these functionalities to the above-defined

20 enantiomerically enriched chiral nitrogen compounds. Particularly preferred functionalities are those which may react with amines to increase the valency of the amine (e.g. form tertiary amines from secondary amines or form secondary amines from primary amines). Examples of such functionalities include chlorine, bromine or iodine atoms, and also perfluoroacylate or sulphonate radicals.

25

The organic modifications described, i.e. the preparative methods for activated supports, are sufficiently well known and may be carried out in a manner known per se, for example, by



reacting the support with a chlorinating agent, for example carbon tetrachloride, thionyl chloride, titanium tetrachloride or phosphorus pentachloride (see also Beck et al. J. Am. Chem. Soc 1992, 114, 10834) and subsequently reacting with functionalized alcohols or alkoxy silanes or

5

- reacting the support with silicon tetrachloride, subsequently reacting with secondary amines and reacting further with functionalized alcohols or alkoxy silanes (see also Petrucci et al. Bull. Chem. Soc. Japan, 1990, 63, 988) silazanes or preferably by


10

- reacting the support with silicon tetrachloride or chlorosilanes of the  $\text{SiCl}_r(\text{C}_1\text{-C}_8\text{-alkyl})_s(\text{C}_4\text{-C}_{14}\text{-Aryl})_t(\text{C}_5\text{-C}_{15}\text{-arylalkyl})_u$  type where  $r$  is one, two or three and  $r+s+t+u = 4$ , and subsequently reacting with functionalized alcohols, chlorosilanes or alkoxy silanes.

15

Particularly preferred **activated supports** are those which are obtainable by reacting supports with chlorosilanes of the  $\text{SiCl}_r(\text{C}_1\text{-C}_8\text{-alkyl})_s(\text{C}_4\text{-C}_{14}\text{-aryl})_t(\text{C}_5\text{-C}_{15}\text{-arylalkyl})_u$  type where  $r$  is one, two or three and  $r+s+t+u = 4$  and subsequently reacting with functionalized alkoxy silanes or halosilanes of the formulae (IIIa) or

20 (IIIb)



25 where, in formula (III),

Hal is chlorine or bromine.

Very particularly preferred **activated supports** are those which are obtained by reacting supports with diphenyldichlorosilane and subsequently reacting with 3-bromopropyltrichlorosilane.


5 Also in formula (I),

(M<sup>m+</sup>) is preferably cobalt in the formal oxidation states 0, +2 and +3, rhodium and iridium in the formal oxidation states +1 and +3, nickel, palladium and platinum in the formal oxidation states 0 and +2 and also ruthenium in the formal oxidation state +2, and  
10 preference is given to Rh<sup>I</sup>, Ir<sup>I</sup> and Pd<sup>II</sup>.

L is preferably the following ligand types: monoolefins, for example ethylene, cyclooctene and cyclohexene, diolefins, for example 1,5-cyclooctadiene (cod), norbornadiene (nbd) and butadiene, nitriles  
15 such as acetonitrile (ACN), benzonitrile and benzylnitrile, aromatics such as benzene, mesitylene and cymene, and also anionic ligands such as allyl, methylallyl, phenylallyl, C<sub>1</sub>-C<sub>8</sub>-alkyl acyl-acetates, C<sub>1</sub>-C<sub>8</sub>-alkyl acylates, chloride, bromide and iodide.

20




(An<sup>q-</sup>) is preferably non-coordinating or weakly coordinating anions, for example nitrate, perchlorate, sulphate, hexafluorophosphate, hexafluoroantimonate, hexachloroantimonate, borates, for example tetrafluoroborate and tetraphenylborate or sulphonates, for example  
25 trifluoromethanesulphonate and nonafluorobutanesulphonate.




as an entire fragment is particularly preferably Rh(cod)BF<sub>4</sub>, Ir(cod)BF<sub>4</sub>, Rh(cod)PF<sub>6</sub>, Ir(cod)PF<sub>6</sub>, Rh(cod)SbF<sub>6</sub>, Ir(cod)SbF<sub>6</sub>, Rh(cod)ClO<sub>4</sub>, Ir(cod)ClO<sub>4</sub>, Rh(nbd)BF<sub>4</sub>, Ir(nbd)BF<sub>4</sub>, Rh(nbd)PF<sub>6</sub>, Ir(nbd)PF<sub>6</sub>, Rh(nbd)SbF<sub>6</sub>, Ir(nbd)SbF<sub>6</sub>, Rh(nbd)ClO<sub>4</sub>, Ir(nbd)ClO<sub>4</sub>, Pd(allyl)BF<sub>4</sub>, Pd(allyl)PF<sub>6</sub> and Pd(ACN)<sub>2</sub>(BF<sub>4</sub>)<sub>2</sub>.

5

Very particularly preferred compounds of the formula (I) are those of the formulae (Ia), (Ib), (Ic) and (Id)





where, in each case,

\* marks a stereogenic centre which is either R- or S-configured, with the proviso that mesoforms are excluded (compounds of the formula (Ic) and (Id))


$M^+$  is rhodium<sup>I</sup> or iridium<sup>I</sup> and

L is cod or nbd and

$An^-$  is perchlorate, hexafluorophosphate, trifluoromethanesulphonate or tetrafluoroborate.


To prepare the compounds of the formula (I), in particular those of the formulae (Ia) to (Id), the procedure is preferably that,

in a step A), an activated support of the formula (IV)




where linker and support have the definitions and areas of preference specified under the formula (I)

are reacted with enantiomerically enriched chiral nitrogen compounds of the formula (IIa)



where the definitions and areas of preference specified under the formula (I) apply, optionally in the presence of an organic solvent, and, in a step B), the compounds of the formula (V) obtained in this way



are reacted with transition metal compounds, optionally in the presence of an organic solvent, to give compounds of the formula (I).

The invention also encompasses the compounds of the formula (V), with the areas of preference specified under the formula (I) applying in the same manner.

Preferred transition metal compounds are those of the formula (VIa)




where

M<sup>1</sup> is ruthenium, rhodium, iridium, nickel, palladium or platinum and

An<sup>1</sup> is chloride, bromide, acetate, nitrate, methanesulphonate, trifluoromethane-sulphonate or acetylacetonate and

5 P<sup>1</sup> for ruthenium, rhodium and iridium is 3, and for nickel, palladium and platinum is 2,

or transition metal compounds of the formula (VIb)



10 where

M<sup>2</sup> is ruthenium, rhodium, iridium, nickel, palladium or platinum and


15 An<sup>2</sup> is chloride, bromide, acetate, methanesulphonate, trifluoromethane-sulphonate, tetrafluoroborate, hexafluorophosphate perchlorate, hexafluoroantimonate, tetra(bis-3,5-trifluoromethylphenyl)borate or tetraphenylborate and

20 p<sub>2</sub> is rhodium and iridium is 1, and for nickel, palladium, platinum and ruthenium is 2 and

L<sup>1</sup> is in each case a C<sub>2</sub>-C<sub>12</sub>-alkene, for example ethylene or cyclooctene, or a nitrile, for example acetonitrile, benzonitrile or benzyl nitrile, or

25 L<sup>1</sup><sub>2</sub> together is a (C<sub>4</sub>-C<sub>12</sub>)-diene, for example norbornadiene or 1,5-cyclooctadiene,

or transition metal compounds of the formula (VIc)



where

$M^3$  is ruthenium and

5

$L^2$  is cod, nbd, allyl, methylallyl or aryl radicals, for example cymene, mesitylene, benzene and

$An^3$  is chloride, bromide, acetate, methanesulphonate, trifluoromethanesulphonate, tetrafluoroborate, hexafluorophosphate perchlorate, hexafluoroantimonate, tetra(bis-3,5-trifluoromethylphenyl)borate or tetraphenylborate,

10

or transition metal compounds of the formula (VIId)

15



where

$M^5$  is palladium, nickel, iridium or rhodium and

20


$An^3$  is chloride or bromide and

$M^4$  is lithium, sodium, potassium, ammonium or organic ammonium and

25

$p^3$  for rhodium and iridium is 3, and for nickel, palladium and platinum is 2,

or transition metal compounds of the formula (VId)



where

$M^6$  is iridium or rhodium and

5  $L^3$  is a ( $C_4$ - $C_{12}$ )-diene, for example norbornadiene or 1,5-cyclooctadiene, and

$An^4$  is a non-coordinating or weakly coordinating anion, for example methane-sulphonate, trifluoromethanesulphonate, tetrafluoroborate, hexafluorophosphate perchlorate, hexafluoroantimonate, tetra(bis-3,5-trifluoromethyl-phenyl)borate or tetraphenylborate.

Examples of further suitable transition metal compounds include  $Ni(cod)_2$ ,  $Pd_2(dibenzylideneacetone)_3$ , cyclopentadienyl $_2Ru$ ,  $Rh(acetylacetonate)(CO)_2$ ,  $Ir(pyridine)_2(cod)OTf$  or multinuclear bridged complexes, for example

15  $[Pd(allyl)Cl]_2$ ,  $[Pd(allyl)Br]_2$ ,  $[Rh(cod)Cl]_2$ ,  $[Rh(cod)Br]_2$ ,  $[Rh(ethene)_2Cl]_2$ ,  $[Rh(cyclooctene)_2Cl]_2$ ,  $[Ir(cod)Cl]_2$  and  $[Ir(cod)Br]_2$ ,  $[Ir(ethene)_2Cl]_2$  and  $[Ir(cyclooctene)_2Cl]_2$ .

Particularly preferred transition metal compounds are:

20  $[Pd(allyl)Cl]_2$ ,  $[Pd(allyl)Br]_2$ ,  $[Rh(cod)Cl]_2$ ,  $[Rh(cod)_2Br]$ ,  $[Rh(cod)_2]ClO_4$ ,  $[Rh(cod)_2]BF_4$ ,  $[Rh(cod)_2]PF_6$ ,  $[Rh(cod)_2]Otf$  ( $Otf$  = triflate),  $[Rh(cod)_2]BPh_4$ ,  $[Rh(cod)_2]SbF_6$ ,  $RuCl_2(cod)$ ,  $[(cymene)RuCl_2]_2$ ,  $[(benzene)RuCl_2]_2$ ,  $[(mesitylene)RuCl_2]_2$ ,  $[(cymene)RuBr_2]_2$ ,  $[(cymene)RuI_2]_2$ ,  $[(cymene)Ru(BF_4)_2]_2$ ,  $[(cymene)Ru(PF_6)_2]_2$ ,  $[(cymene)Ru(BPh_4)_2]_2$ ,  $[(cymene)Ru(SbF_6)_2]_2$ ,  $[Ir(cod)_2Cl]_2$ ,  $[Ir(cod)_2]PF_6$ ,

25  $[Ir(cod)_2]ClO_4$ ,  $[Ir(cod)_2]SbF_6$ ,  $[Ir(cod)_2]BF_4$ ,  $[Ir(cod)_2]Otf$ ,  $[Ir(cod)_2]BPh_4$ ,  $[Rh(nbd)Cl]_2$  ( $nbd$  = norbornadiene),  $[Rh(nbd)_2Br]$ ,  $[Rh(nbd)_2]ClO_4$ ,  $[Rh(nbd)_2]BF_4$ ,  $[Rh(nbd)_2]PF_6$ ,  $[Rh(nbd)_2]Otf$ ,  $[Rh(nbd)_2]BPh_4$ ,  $[Rh(nbd)_2]SbF_6$ ,  $RuCl_2(nbd)$ ,  $[Ir(nbd)_2]PF_6$ ,  $[Ir(nbd)_2]ClO_4$ ,  $[Ir(nbd)_2]SbF_6$ ,  $[Ir(nbd)_2]BF_4$ ,  $[Ir(nbd)_2]Otf$ ,  $[Ir(nbd)_2]BPh_4$ ,  $Ir(pyridine)_2(nbd)Otf$ ,  $[Ru(DMSO)_4Cl_2]$ ,



[Ru(ACN)<sub>4</sub>Cl<sub>2</sub>], [Ru(PhCN)<sub>4</sub>Cl<sub>2</sub>] and [Ru(cod)Cl<sub>2</sub>]<sub>n</sub>, and even greater preference is given to [Pd(allyl)Cl]<sub>2</sub>, Rh(cod)<sub>2</sub>OTf, Rh(cod)<sub>2</sub>PF<sub>6</sub>, Rh(cod)<sub>2</sub>SbF<sub>6</sub>, Ir(cod)<sub>2</sub>BF<sub>4</sub>, Rh(cod)<sub>2</sub>OTf, Ir(cod)<sub>2</sub>PF<sub>6</sub>, Ir(cod)<sub>2</sub>SbF<sub>6</sub> and Ir(cod)<sub>2</sub>BF<sub>4</sub>.

- 5 It is pointed out that it is often advantageous when using halide-containing transition metal compounds to additionally use silver or potassium salts of non-coordinating or weakly coordinating anions as defined above in an approximately equimolar amount to the halide present.
- 10 Useful organic solvents for steps A) and B) are typically aliphatic or aromatic, optionally halogenated hydrocarbons, for example petroleum ether, benzene, toluene, the isomeric xylenes, chlorobenzene, the isomeric dichlorobenzenes, hexane, cyclohexane, dichloromethane or chloroform, and preferably ethers, such as diethyl ether, diisopropyl ether, dioxane, tetrahydrofuran, methyl tert-butyl
- 15 ether or ethylene glycol dimethyl ether or ethylene glycol diethyl ether. Particularly preferred organic solvents are toluene, diethyl ether, tetrahydrofuran and methyl tert-butyl ether.

- The weight ratio of enantiomerically enriched chiral nitrogen compounds to
- 20 activated support may be, for example and with preference, 0.02:1 to 100:1, particularly preferably 0.1:1 to 5:1 and very particularly preferably 0.1:1 to 1:1. The use of greater amounts of enantiomerically enriched chiral nitrogen compounds is possible but uneconomic.

- 25 The weight ratio of transition metal compound to compounds of the formula (V) may be, for example and with preference, 0.005:1 to 1:1, particularly preferably 0.01:1 to 0.2:1 and very particularly preferably 0.02:1 to 0.1:1.

In step A), the reaction temperature is, for example and with preference, 0 to 100°C, particularly preferably 20 to 80°C and very particularly preferably 40 to 60°C.

- 5 In step B), the reaction temperature is, for example and with preference, 20°C to 80°C, particularly preferably 0 to 60°C and very particularly preferably 10 to 30°C.

- 10 The compounds of the formula (I) may be worked up in a manner known per se by filtration and/or centrifugation and/or sedimentation and optionally subsequent washing with organic solvent, and the washing may be carried out, for example, batchwise or continuously. For storage purposes, the compounds of the formula (I) are preferably dried.

- 15 The compounds of the formula (I) may be used directly as catalyst for asymmetric reactions.

The invention therefore also encompasses catalysts which comprise compounds of the formula (I).


20

The invention also encompasses a process for catalytically preparing enantiomerically enriched compounds, which is characterized in that the catalysts used are those which comprise compounds of the formula (I).

- 25 Preferred processes for preparing enantiomerically enriched compounds are asymmetric hydrogenations, for example hydrogenations of prochiral C=C bonds such as prochiral enamines, olefins, enol ethers; C=O bonds such as prochiral ketones and C=N bonds such as prochiral imines. Particularly preferred

asymmetric hydrogenations are hydrogenations of prochiral ketones, in particular alpha- and beta-ketocarboxylic esters.

Preferred alpha- and beta-ketocarboxylic esters are compounds of the formula  
5 (VII)



where

10  $\text{R}^5$  and  $\text{R}^7$  are each independently  $\text{C}_1$ - $\text{C}_{12}$ -alkyl,  $\text{C}_1$ - $\text{C}_{12}$ -haloalkyl,  $\text{C}_5$ - $\text{C}_{15}$ -aryl-alkyl or  $\text{C}_4$ - $\text{C}_{14}$ -aryl and

$\text{R}^6$  is absent or is 1,1-( $\text{C}_1$ - $\text{C}_4$ -alkylene).

15 Preferably,  $\text{R}^5$  and  $\text{R}^7$  are each independently  $\text{C}_1$ - $\text{C}_4$ -alkyl or phenyl, and  $\text{R}^6$  is methylene or is absent.

Particularly preferred compounds of the formula (VII) are methyl phenyl-glyoxylate and ethyl chloroacetoacetate.

20

The hydrogenation according to the invention of alpha- and beta-ketocarboxylic esters provides enantiomerically enriched compounds of the formula (VIII)



where

\* marks a stereogenic centre which is S- or R-configured and

- 5     $R^5$ ,  $R^6$  and  $R^7$  each have the definitions and areas of preference specified under the formula (VII).

10    The compounds which can be prepared according to the invention are suitable in particular as optical resolution reagents or in a process for preparing pharmaceuticals or agrochemicals.

15    In a preferred embodiment of asymmetric hydrogenations according to the invention, the reaction temperature is 0 to 200°C, preferably 10 to 150°C, and the partial hydrogen pressure is, for example, 0.1 to 200 bar, preferably 0.9 to 100 bar and particularly preferably 4 to 30 bar.

20    Useful solvents for asymmetric hydrogenations according to the invention are in particular aliphatic or aromatic, optionally halogenated hydrocarbons, for example petroleum ether, benzene, toluene, the isomeric xylenes, chlorobenzene, the isomeric dichlorobenzenes, hexane, cyclohexane, dichloromethane or chloroform, ethers such as diethyl ether, diisopropyl ether, dioxane, tetrahydrofuran, methyl tert-butyl ether or ethylene glycol dimethyl ether or ethylene glycol diethyl ether, and preferably alcohols such as methanol, ethanol and isopropanol.

25    The weight ratio of compounds of the formula (I) to substrate may be, for example, 1:1 to 1:10.000, preferably 1:5 to 1:1000.

The advantage of the present invention is that heterogeneous catalysts may be prepared in high yields and in an efficient manner and that these catalysts allow

high conversions and enantioselectivities in asymmetric syntheses. This fact is to be regarded as particularly surprising in that the non-immobilized analogues to compounds of the formula (I) allow only very low enantioselectivities, if any at all, as the comparative examples show.

5

### Examples

#### Example 1 (for comparison)

#### 10 Preparation of [palladium( $\eta^3$ -allyl)((S)-2-aminomethyl-1-ethylpyrrolidine)]-BF<sub>4</sub>

[Pd( $\eta^3$ -allyl)Cl]<sub>2</sub> (70 mg, 0.19 mmol) were dissolved in THF (10 ml), admixed with AgBF<sub>4</sub> (80 mg, 0.41 mmol) and stirred for one hour. The mixture was  
15 filtered, the amine (48 mg, 0.38 mmol) was added to the filtrate and the mixture was stirred for 30 min. A white solid precipitated out. The further addition of 20 ml of hexane resulted in further product precipitating out. The solution was filtered, washed with hexane (2 x 20 ml) and diethyl ether (2 x 20 ml) and the residue was dried under reduced pressure to obtain a white powder (100 mg, 73%  
20 of theory).

Anal: Calculated: C<sub>10</sub>H<sub>21</sub>N<sub>2</sub>PdBF<sub>4</sub>: C, 33.14; H, 5.80; N, 7.73. Found: C, 33.40; H, 5.81; N, 7.55.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): 1.30-4.0 (m, CH<sub>2</sub> allyl and amine, 18H) 5.55 (m, CH, 1H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): 14.8(1), 22.2(4), 25.3(5), 44.2(7), 50.1(2), 53.2(3), 60.3 (CH<sub>2</sub>  
25 allyl), 69.9(6) 116.3(CH, allyl).

ESI = 275 (M<sup>+</sup>), 233 (M<sup>+</sup> - allyl)

## **Example 2**

### **Preparation of covalently immobilized [palladium( $\eta^3$ -allyl)((S)-2-aminomethyl-1-ethylpyrrolidine)]BF<sub>4</sub>**

5

#### **a) Activation of MCM-41**

Dichlorodiphenylsilane (0.48 g) was added to dried, calcined MCM-41 (2.0 g) in THF (15 ml) and stirred for one hour. The solution was then cooled to  $-78^{\circ}\text{C}$  and admixed with 3-bromopropyltrichlorosilane (1.10 g). The mixture was allowed to  
10 warm slowly to room temperature and stirred for a further 8 hours. The mixture was subsequently stirred at  $50^{\circ}\text{C}$  for one hour. The activated support MCM-41 was filtered off and purified by Soxhlet extraction using THF. Finally, the support was dried under reduced pressure.

Anal: C, 4.70; H, 0.87.

15

#### **b) Coupling with (S)-2-aminomethyl-1-ethylpyrrolidine**

The activated support MCM-41 (700 mg) in THF (15 ml) together with the amine (0.15 ml) was heated at  $50^{\circ}\text{C}$  for 20 hours. Afterwards, the product was filtered and purified by Soxhlet extraction using THF. Finally, the product was dried  
20 under reduced pressure.

Anal: C, 9.82; H, 2.0; N, 1.51.

#### **c) Complexing with palladium**

In a Schlenk vessel, [PdCl(allyl)]<sub>2</sub> (30 mg, 0.08 mmol) and AgBF<sub>4</sub> (30 mg, 0.15 mmol) were dissolved in THF (10 ml) and stirred for one hour. The solution  
25 was filtered, admixed with the modified MCM-41 from b) and stirred for four hours. The white solid was filtered, washed with THF (4 x 20 ml) and dried under reduced pressure.

Anal: C, 11.67; H, 2.32; N, 1.46.

**Example 3 (for comparison)****Preparation of [palladium( $\eta^3$ -allyl)((1R,2R)-1,2-diphenylethylenediamine) ]BF<sub>4</sub>**

5. [Pd( $\eta^3$ -allyl)Cl]<sub>2</sub> (70 mg, 0.19 mmol) were dissolved in THF (10 ml), admixed with AgBF<sub>4</sub> (80 mg, 0.41 mmol) and stirred for one hour. The mixture was filtered, the amine (80 mg, 0.38 mmol) was added to the filtrate and the mixture was stirred for 30 min. A white solid precipitated out. The further addition of
- 10 20 ml of hexane resulted in further product precipitating out. The solution was filtered, washed with hexane (2 x 20 ml) and diethyl ether (2 x 20 ml) and the residue was dried under reduced pressure to obtain a white powder (156 mg, 88% of theory).
- Anal.: Calculated for C<sub>17</sub>H<sub>21</sub>N<sub>2</sub>RhBF<sub>4</sub>: C, 45.74; H, 4.71; N, 6.27. Found: C,
- 15 45.25; H, 4.62; N, 5.98.
- <sup>1</sup>H NMR (CD<sub>3</sub>OD) 2.98 (m, CH<sub>anti</sub>, 2H), 4.02 (s, NCH, 2H), 4.20 (m, CH<sub>syn</sub>, 2H), 5.47 (m, CH<sub>central</sub>, 1H), 7.1-7.25 (m, Ph, 10H).
- <sup>13</sup>C NMR (CD<sub>3</sub>OD) 56.72 (CH<sub>2</sub>), 64.24 (NCH), 114.87 (CH), 127.0, 127.70, 128.25, 139.73 (Ph).
- 20 +ve ESI = 359 (M<sup>+</sup>).

**Example 4**

- Preparation of covalently immobilized [palladium( $\eta^3$ -allyl)((1R,2R)-1,2-**
- 25 **diphenylethylenediamine) ]BF<sub>4</sub>**

**a) Activation of MCM-41**

Dichlorodiphenylsilane (0.48 g) was added to dried, calcined MCM-41 (2.0 g) in THF (15 ml) and stirred for one hour. The solution was then cooled to -78°C and

admixed with 3-bromopropyltrichlorosilane (1.10 g). The mixture was allowed to warm slowly to room temperature and stirred for a further 8 hours. The mixture was subsequently stirred at 50°C one hour for one hour. The activated support MCM-41 was filtered and purified by Soxhlet extraction using THF. Finally, the support was dried under reduced pressure.

Anal: C, 4.70; H, 0.87.

**b) Coupling with (1R,2R)-1,2-diphenylethylenediamine**

The activated support MCM-41 (500 mg) in THF (15 ml) together with the amine (150 mg) was heated to reflux for 20 hours. Afterwards, the product was filtered and purified by Soxhlet extraction using THF. Finally, the product was dried under reduced pressure.

Anal: C, 9.72; H, 1.92; N, 0.15.

**c) Complexing with palladium**

In a Schlenk vessel, [PdCl(allyl)]<sub>2</sub> (55 mg, 0.15 mmol) and AgBF<sub>4</sub> (60 mg, 0.3 mmol) were dissolved in THF (10 ml) and stirred for one hour. The solution was filtered, admixed with the modified MCM-41 from b) and stirred for four hours. The solid was filtered, washed with THF (4 x 20 ml) and dried under reduced pressure.

Anal: C, 10.67; H, 2.31; N, 0.13.

**Example 5 (for comparison)**

**Preparation of [Rh(cod)((1R,2R)-1,2-diphenylethylenediamine)]BF<sub>4</sub>**

[RhCl(cod)]<sub>2</sub> (64 mg, 0.13 mmol) was dissolved in THF (10 ml), AgCF<sub>3</sub>SO<sub>3</sub> (50 mg, 0.26 mmol) was added and the solution was stirred for one hour. The solution was subsequently filtered, the filtrate admixed with (1R, 2R)-1,2-



diphenylethylenediamine (50 mg, 0.26 mmol) and the resulting solution was stirred for one hour. Subsequently, the solution was concentrated under reduced pressure and admixed with hexane (25 ml), and the product precipitated out. The mixture was filtered, and the product was washed with hexane (2 x 20 ml) and diethyl ether (2 x 20 ml) and dried under reduced pressure. A yellow powder was obtained (185mg, 90%).

Anal: Calculated for  $C_{22}H_{28}N_2RhBF_4$  C, 51.76; H, 5.50; N, 5.50. Found: C, 51.44; H, 5.57; N, 5.29.

$^1H$  NMR ( $CD_3OD$ ) 1.95 (br m,  $CH_2$  4H), 2.46 (br m,  $CH_2$  4H), 4.01 (s,  $NCH$ , 2H), 4.24 (m,  $CH$ , 2H), 4.35 (m,  $CH$ , 2H), 7.1-7.3 (m, Ph, 10H).

$^{13}C$  NMR ( $CD_3OD$ ) 31.6 ( $CH_2$ ), 66.3 ( $NCH$ ) 81.4 ( $CH$ ), 128.5, 129.2, 129.6, 140.5 (Ph).

+ve ESI = 423 ( $M^+$ ).

## 15 **Example 6**

**Preparation of covalently immobilized  $[Rh(cod)((1R,2R)\text{-}1,2\text{-diphenylethylene-diamine})]BF_4$**

### 20 **a) Activation of MCM-41**

Dichlorodiphenylsilane (0.48 g) was added to dried, calcined MCM-41 (2.0 g) in THF (15 ml) and stirred for one hour. The solution was then cooled to  $-78^\circ C$  and admixed with 3-bromopropyltrichlorosilane (1.10 g). The mixture was allowed to warm slowly to room temperature and stirred for a further 8 hours. The mixture was subsequently stirred at  $50^\circ C$  one hour for one hour. The activated support MCM-41 was filtered and purified by Soxhlet extraction using THF. Finally, the support was dried under reduced pressure.

Anal: C, 4.70; H, 0.87.

**b) Coupling with (1R,2R)-1,2-diphenylethylenediamine**

The activated support MCM-41 (500 mg) in THF (15 ml) together with the amine (150 mg) was heated to reflux for 24 hours. Afterwards, the product was filtered and purified by Soxhlet extraction using THF. Finally, the product was dried  
5 under reduced pressure.

Anal: C, 9.72; H, 1.92; N, 0.15; Br, 1.30.

**c) Complexing**

In a Schlenk vessel,  $[\text{RhCl}(\text{cod})]_2$  (30 mg, 0.06 mmol) and  $\text{AgBF}_4$  (30 mg, 0.15 mmol) were dissolved in THF (10 ml) and stirred for one hour. The solution  
10 was filtered, admixed with the modified MCM-41 from b) and stirred for four hours. The solution decolorized slowly. The yellow solid was filtered, washed with THF (4 x 20 ml) and dried under reduced pressure.

Anal: C, 11.23; H, 2.02; N, 0.12.

15

**Example 7 (for comparison)**

**Preparation of  $[\text{Rh}(\text{cod})((\text{S})\text{-2-aminomethyl-1-ethylpyrrolidine})]\text{BF}_4$**

$[\text{RhCl}(\text{cod})]_2$  (100 mg, 0.20 mmol) was dissolved in THF (10 ml),  $\text{AgCF}_3\text{SO}_3$   
20 (80 mg, 0.41 mmol) was added and the solution was stirred for one hour. The solution was subsequently filtered, the filtrate admixed with (S)-2-aminomethyl-1-ethylpyrrolidine (51.2 mg, 0.40 mmol) and the resulting solution was stirred for one hour. Subsequently, the solution was concentrated under reduced pressure and admixed with hexane (25 ml), and the product precipitated out. The mixture was  
25 filtered, and the product was washed with hexane (2 x 20 ml) and diethyl ether (2 x 20 ml) and dried under reduced pressure. A yellow powder was obtained (185 mg, 90%).

Anal.: Calculated for  $\text{C}_{15}\text{H}_{28}\text{N}_2\text{RhBF}_4$ : C, 42.25; H, 6.57; N, 6.57. Found: C, 42.79; H, 6.61; N, 6.59.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 1.76 (br m, 4H,  $\text{CH}_2$  olefin), 2.45 (br m, 4H,  $\text{CH}_2$  olefin), 1.55-3.30 (m, 14H, amine).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) = 12.3 (1), 21.7 (4), 24.3 (5), 45.9 (7), 51.0 (2), 56.6 (3), 70.0 (6), 30.4, 30.7 ( $\text{CH}_2$ ) 79.7, 83.6 (CH).

5 +ve ESI = 339 ( $\text{M}^+$ ), 231 ( $\text{M}^+ - \text{COD}$ ).

### Example 8

10 **Preparation of covalently immobilized  $[\text{Rh}(\text{cod})((\text{S})\text{-2-aminomethyl-1-ethylpyrrolidine})\text{BF}_4$**

#### **a) Activation of MCM-41**

Dichlorodiphenylsilane (0.48 g) was added to dried, calcined MCM-41 (2.0 g) in THF (15 ml) and stirred for one hour. The solution was then cooled to  $-78^\circ\text{C}$  and  
15 admixed with 3-bromopropyltrichlorosilane (1.10 g). The mixture was allowed to warm slowly to room temperature and stirred for a further 8 hours. The mixture was subsequently stirred at  $50^\circ\text{C}$  one hour for one hour. The activated support MCM-41 was filtered and purified by Soxhlet extraction using THF. Finally, the support was dried under reduced pressure.

20 Anal: C, 4.70; H, 0.87.

#### **b) Coupling with (S)-2-aminomethyl-1-ethylpyrrolidine**

The activated support MCM-41 (700 mg) in THF (15 ml) together with the amine (0.15 ml) was heated at  $50^\circ\text{C}$  for 20 hours. Afterwards, the product was filtered  
25 and purified by Soxhlet extraction using THF. Finally, the product was dried under reduced pressure.

Anal: C, 9.82; H, 2.0; N, 1.51.

**c) Complexing with palladium**

In a Schlenk vessel,  $[\text{RhCl}(\text{cod})]_2$  (30 mg, 0.06 mmol) and  $\text{AgBF}_4$  (30 mg, 0.15 mmol) were dissolved in THF (10 ml) and stirred for one hour. The solution was filtered, admixed with the modified MCM-41 from b) and stirred for four  
5 hours. The white solid was filtered, washed with THF (4 x 20 ml) and dried under reduced pressure.

Anal: C, 12.45; H, 2.45; N, 1.60.

The  $^{19}\text{F}$  MAS NMR spectrum confirmed the presence of the  $\text{BF}_4^-$  anion.

10 **Examples 9 to 15**

**General procedure for the use of the catalysts in asymmetric hydrogenations**

The asymmetric hydrogenations were carried out in a high-pressure autoclave  
15 made of rust-free stainless steel and having a capacity of 150 ml. 10 mg in each case of the homogeneous catalyst or 50 mg in each case of the immobilized catalysts were transferred into the high-pressure autoclave under an inert atmosphere.

20 Methyl phenylglyoxylate (0.5 g), methanol (30 g) and an internal standard (cyclo-dodecane) were added and the high-pressure autoclave was closed. The high-pressure autoclave and its inlets and outlets were subsequently inertized by flushing with nitrogen three times and, to test the seal, finally placed under a hydrogen pressure of 5 bar. Subsequently, the hydrogen pressure was increased to  
25 20 bar, the high-pressure autoclave was brought to reaction temperature (313 K) and the contents were stirred with a mechanical stirrer at 400 rpm.

A miniaturized automatic withdrawal valve was used to take samples of the contents, in order to be able to investigate the progress of the reaction. At the end

of the reaction, the high-pressure autoclave was cooled for two hours in an ice bath and decompressed, and the products were identified by gas chromatography (GC, Varian, Model 3400 CX) using a chiral column (Chiraldex, 20 m x 0.25 mm).

5

The results of the hydrogenation experiments are compiled in the following table:

Example	Catalyst from example	Temperature [°C]	Reaction time [h]	Conversion [%]	ee [%]
9	1 (for comp.)	40	2	97.3	0
10	1 for comp.)	40	24	99.0	0
11	2	40	2	99.1	94.5
12	5 (for comp.)	40	2	82.0	0
13	7 (for comp.)	40	2	94.0	0
14	8	40	2	97.7	77.5
15	8	40	24	98.6	80.4

10 Although the invention has been described in detail in the foregoing for the purpose of illustration, it is to be understood that such detail is solely for that purpose and that variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention except as it may be limited by the claims.